I N D E X

S. No.	Contents	Page No.
1.	**SET THEORY & COUNTABILITY**	**1**
1.1	Sets	1
1.2	Relation	2
1.3	Function	3
1.4	Classification of Functions	4
1.5	Similar/Equivalent Sets	7
1.6	Countable & Uncountable Sets	8
1.7	Inequality	12
2.	**POINT SET TOPOLOGY**	**16**
2.1	Archimedean Property	16
2.2	Bounds of A Set	16
2.3	Intervals	18
2.4	Connectedness	18
2.5	Neighbourhood of A Point	18
2.6	Some Important Type of Points & Its Properties	19
3.	**SEQUENCES**	**29**
3.1	Definitions	29
3.2	Bounded & Monotonic Sequences	29
3.3	Subsequence's of A Sequence	31
3.4	Limit of A Sequence	33
3.5	Non Convergent Sequence	39
3.6	Some Important Limits	42
3.7	Limit Superior & Limit Inferior	49
3.8	Sequence of Natural Number's	54
3.9	Sequence of Rational Numbers	55
3.10	Cauchy Sequences	55

4.	**SERIES OF REAL NUMBERS**		**64**
	4.1	Sequence of Partial Sum	64
	4.2	Positive Term Series	65
	4.3	Some Important Series & Their Convergence	67
	4.4	Important Test for Convergence	70
	4.5	Alternating Term Series	77
	4.6	Series of Arbitrary Terms	82
	4.7	Sum of Series	84
5.	**ELEMENTARY FUNCTIONS**		**97**
	5.1	Function	97
	5.2	Wavy Curve Method	98
	5.3	Some Well Known Functions	99
	5.4	Trigonometric Functions	100
	5.5	Inverse Trignometric Function	101
	5.6	Graphical Transformations	102
	5.7	Algebraic & Transcendental Function	105
	5.8	Bounded Function	106
6.	**LIMIT & CONTINUITY**		**109**
	6.1	Limit of A Function	109
	6.2	Left Hand Limit & Right Hand Limit	111
	6.3	Continuity	114
	6.4	Discontinuity	115
	6.5	Cauchy Functional Equation	117
	6.6	Properties of Continuous Functions	118
	6.7	Roots of A Function	121
	6.8	Lipschitz Function	124
	6.9	Uniform Continuity	124

7. DIFFERENTIABILITY 138

7.1 Differentiability..... 138

7.2 Monotonic Functions & Differentiability.....143

7.3 Fixed Point..... 144

7.4 Local Extrema of a Function.....146

7.5 Application of Differentiability..... 147

8. RIEMANN INTEGRATION & BOUNDED VARIATION 159

8.1 Riemann Integration..... 159

8.2 Definite Integral as Limit of Sum..... 161

9. IMPROPER INTEGRAL 166

10. FUNCTION OF SEVERAL VARIABLE 171

10.1 Some Important Definitions..... 171

10.2 Limit of A Function..... 172

10.3 Continuity..... 176

10.4 Directional Derivatives.....178

10.5 Differentiability..... 180

10.6 Partial Derivatives of Higher Order 183

10.7 Extreme Values..... 184

10.8 Function of n Variables.....186

10.9 Vector Valued Function..... 186

CHAPTER 1

SET THEORY & COUNTABILITY

SET THEORY

1.1 SETS :

1.1.1 DEFINITION:

A Set is a well-defined collection of distinct objects.

The individual objects of the set are called member or elements of the set.

Set is denoted by Capital letters A, B, C, X, Y, Z ...

Examples :-

(1) Set of all vowels in English Alphabet $X = \{a, e, i, o, u\}$.

(2) Some Important sets : $\mathbb{N}$ = set of all natural numbers

$\mathbb{Z}$ = set of all integers

$\mathbb{R}$ = set of all real numbers

$\mathbb{C}$ = set of all complex numbers

$\mathbb{Q}$ = set of all rational numbers

$\mathbb{Q}^c$ = set of all irrational numbers

(3) The set of all natural numbers which are strictly less than 4.

$X = \{x \in \mathbb{N} \mid x < 4\} = \{1,2,3\}$

Note:

(1) Set guarantees that the Collection will contains distinct elements but collection doesn't guarantee this.

(2) The word 'Well-defined' means there is no ambiguity or confusion as to whether an object belongs to collection or not.

1.1.2 Equality of Two sets :

Two sets are said to be equal when they consist of exactly the same elements.

(i) $\{a, b, c\} = \{b, c, a\} = \{c, b, a\}$

(ii) $\{x \in \mathbb{N} \mid x < 4\} = \{x \in \mathbb{N} | x \leq 3\} = \{1,2,3\}$

Cardinality of the set :

The number of elements in the set A is called cardinality of the set A

Cardinality of the set A is denoted by $|A|$.

Russel's Paradox :

Let R be the collection of all sets that are not members of themselves.

$R = \{A | A \notin A\}$

Exercise: Show that $R \notin R \Leftrightarrow R \in R$.

1.1.3 Subset :

If A & B are two sets such that each member of A is also a member of B

i.e. $x \in A \Rightarrow x \in B$

Then A is called subset of B.

OR,

$\nexists\ x \in A$ such that $x \notin B$.

Notation:

(1) If A is subset of B, then it is denoted by $A \subseteq B$(B is superset of A)

(2) If A is proper subset of B, then it is denoted by $A \subsetneq B$ (i.e. $\exists\ x \in B$ such that $x \notin A$).

Example :

1) If $x \in A \Rightarrow x \in A$

$\Rightarrow A$ is subset of A i.e $A \subseteq A$

$\therefore$ Every set is a subset of itself.

2) ϕ (empty set) is a subset of X i.e. $\phi \subseteq X$.

1.1.4 Power Set :

The collection of all subsets of a set A is called Power set of A.

It is denoted by $P(A)$.

Example :

(1) $A = \{1,2,3\}$

$P(A) = \{\phi, A, \{1\}, \{2\}, \{3\}, \{1,2\}, \{1,3\}, \{2,3\}\}$.

Note:- for any set A

(1) $\phi \in P(A)$

(2) ϕ is element as well as subset of $P(A)$.

1.1.5 Cartesian Product :

Let A & B are two sets. Then Cartesian product of A & B is denoted by $A \times B$ and it is defined as

$A \times B = \{(a, b) | a \in A, b \in B\}$, where (a, b) is an order pair.

Example:-

(1) $A = \{1,2\}, B = \{3,4\}$

$A \times B = \{(1,3), (1,4), (2,3), (2,4)\}$

(2) $A = \{1,2\}, B = \phi$, then

$A \times B = \phi$

1.1.6 Union & Intersection of two sets :

(1) Union :

If A & B are two sets, then the set consisting of all elements which belongs to either A or B is defined as union of A & B.

It is denoted by $A \cup B$

$A \cup B = \{x \mid x \in A \text{ or } x \in B\}$

(2) Intersection :

If A & B are two sets, then the set consisting of all elements which belongs to both A & B is defined as intersection of A & B

It is denoted by $A \cap B$

$A \cap B = \{x | \ x \in A \ and \ x \in B\}$

Note:

(1) Two Set A & B are said to be disjoint if they have no common elements.

i.e. $A \cap B = \phi$

(2) $A \cup \phi = A$, $A \cup A = A$, $A \cup B = B \cup A$

$A \cap \phi = \phi$, $A \cap A = A$, $A \cap B = B \cap A$.

1.1.7 Union & Intersection of an arbitrary family of sets :

Index Set :

An index set is a set whose members labels (or index) members of another set. It is denoted by Λ

Arbitrary Union & Intersection

Let Λ be an index set, then

$\bigcup_{i \in \Lambda} A_n = \{x | \ x \in A_i, \text{ for some } i \in \Lambda\}$

and

$\bigcap_{i \in \Lambda} A_n = \{x | \ x \in A_i, \text{ for each } i \in \Lambda\}$

1.2 Relations :

1.2.1 Definition:

If A & B be two non-empty sets, then a relation from A to B is a subset of $A \times B$ & every subset of $A \times B$ is a relation from A to B

A relation from A to A is called binary relation.

Examples :

1) Null relation $\phi \subseteq A \times B$

2) $A = \{1, 2\}$, & $B = \{0, 1, 2\}$, then

$R_1 = \{(1, 1), (2, 2)\}$ is a relation.

$R_2 = \{(1, 0), (1, 2), (2, 1), (2, 2)\}$ is a relation

$R_3 = \{(1, 0), (0, 1)\}$ is not a relation.

1) Reflexive Relation :

Let R be a binary relation on a set A i.e. $R \subseteq A \times A$.

Then R is called a reflexive relation if

$(a, a) \in R, \forall \ a \in A$

Example :

$A_1 = \{1, 2, 3\}$ & $R_1 = \{(1, 1), (2, 2), (3, 3), (1, 2), (2, 3)\}$

$A_2 = \{a, b\}$ & $R_2 = \{(a, a), (b, b), (a, b), (b, a)\}$

Both are reflexive relation.

2) Symmetric Relation:

Let R be a binary relation on a set A i.e. $R \subseteq A \times A$.

Then R is called a symmetric relation if

$(a, b) \in R \Rightarrow (b, a) \in R$

Example :

$A = \{1, 2, 3\}$ & $R = \{(1, 1), (1, 2), (2, 1), (1, 3), (3, 1)\}$ is symmetric relation.

3) Transitive Relation :

Let R be a binary relation on a set A i.e. $R \subseteq A \times A$.

Then R is called a transitive relation if

(a, b) & $(b, c) \in R$ then $(a, c) \in R$

Example :

$A_1 = \{1, 2, 3\}$, $R_1 = \{(1, 1), (1, 2), (2, 3), (1, 3)\}$ is transitive relation.

Equivalence Relation :

A binary relation $R \subseteq A \times A$ is called equivalence relation if it is reflexive, symmetric & transitive.

1.2.2 Equivalence class :

If $R \subseteq A \times A$ is an equivalence relation. Then equivalence class of A with respect to R is denoted by $[a]$ & is defined as

$[a] = \{x \in A | \ (a, x) \in R\}$

Result :-

(1) Two equivalence class are either equal or disjoint.

(2) Fundamental Theorems of equivalence relation.
An equivalence relation on a set A form a partition of A conversely, Corresponding to any partition of A, there exist an equivalence relation on A.

(3) The number of equivalence relation on a set A is equals to the number of partition of a set A.

1.2.3 Number of Equivalence relations :

Let A be any set with cardinality $|A|$, then

$\|A\|$	Number of equivalence relation
0	1
1	1
2	2
3	5
4	15
5	52

Definition :

Let $S = \phi$, & $R \subseteq S \times S$ be a relation on S, then R is null relation & it is an equivalence relation.

Anti-symmetric Relation:

A binary relation R is anti-symmetric relation if

$(a, b) \in R$ & $(b, a) \in R \Rightarrow a = b$

1.2.4 Partial order Relation :

A relation R on a set A is said to be partial order relation if it is

(a) Reflexive

(b) Anti symmetric

(c) Transitive

Example:

Let $x, y \in \mathbb{R}$, define xRy iff $x - y + \sqrt{2} \in \mathbb{Q}^c$. Then R is

(a) Reflexive relation

(b) Symmetric relation

(c) Transitive relation

(d) Equivalence relation

(e) Anti-symmetric relation

(f) Partial order relation

Solution:

(a) If $y = x$ then $x - y + \sqrt{2} = x - x + \sqrt{2} = \sqrt{2} \in \mathbb{Q}^c$

$\Rightarrow x - y + \sqrt{2} \in \mathbb{Q}^c$

$\Rightarrow x\ R\ x$

$\Rightarrow R$ is a reflexive relation.

(b) $x = 1 + \sqrt{2}, y = 1$

$x - y + \sqrt{2} = 1 + \sqrt{2} - 1 + \sqrt{2} = 2\sqrt{2} \in \mathbb{Q}^c$

$\Rightarrow xRy$

Consider $y - x + \sqrt{2} = 1 - (1 + \sqrt{2}) + \sqrt{2} = 1 - 1 - \sqrt{2} + \sqrt{2} = 0 \notin \mathbb{Q}^c$

$\Rightarrow y$ is not related to x

$\Rightarrow R$ is not a symmetric relation.

(c) $x = \frac{-\sqrt{2}}{2}, y = 1, z = \frac{\sqrt{2}}{2}$

$x - y + \sqrt{2} = \frac{-\sqrt{2}}{2} - 1 + \sqrt{2} = \frac{-\sqrt{2} - 2 + 2\sqrt{2}}{2} = \frac{-2 + \sqrt{2}}{2} \in \mathbb{Q}^c$

$y - z + \sqrt{2} = 1 - \frac{\sqrt{2}}{2} + \sqrt{2} = \frac{2 - \sqrt{2} + 2\sqrt{2}}{2} = \frac{2 + \sqrt{2}}{2} \in \mathbb{Q}^c$

$\Rightarrow xRy\ \&\ yRz$

But $x - z + \sqrt{2} = \frac{-\sqrt{2}}{2} - \frac{\sqrt{2}}{2} + \sqrt{2} = -\frac{2\sqrt{2}}{2} + \sqrt{2} = -\sqrt{2} + \sqrt{2} = 0 \notin \mathbb{Q}^c$

$\Rightarrow R$ is not a transitive relation

(d) Clearly R is not an equivalence relation

(e) $x = \sqrt{2}, \quad y = \sqrt{3}$

$x - y + \sqrt{2} = \sqrt{2} - \sqrt{3} + \sqrt{2} = 2\sqrt{2} - \sqrt{3} \in \mathbb{Q}^c$

$\Rightarrow xRy$

$y - x + \sqrt{2} = \sqrt{3} - \sqrt{2} + \sqrt{2} = \sqrt{3} \in \mathbb{Q}^c$

$\Rightarrow yRx$

But $x \neq y$

$\Rightarrow R$ is not an anti-symmetric relation

(f) Clearly, R is not a partial order relation.

EXERCISE

Let $x, y \in \mathbb{Z}$. Define xRy iff $x = y \pmod 3$. Then R is

(a) Reflexive relation

(b) Symmetric relation

(c) Transitive relation

(d) Equivalence relation

(e) Anti-symmetric relation

(f) Partial order relation

(If R is an equivalence relation, find the equivalence classes)

1.3 Function:

1.3.1 Definition:

Let A & B be any two non-empty sets. Then a rule which assign each member x of A to unique member y of B is called a function and y is the image of x under f.

Notation:

$f : A \to B$ i.e. f is a mapping or a function from A to B.

Note :

(i) A function is a special type of relation in which each element of A is pair with exactly one element of B.

i.e. if $(a, b) \in f \Rightarrow f(a) = b$,

1.3.2 Definition:


Domain, Co-domain & Range of :

Let $f : A \to B$ be any function from A to B then

i) The set A is called the domain of the function f.

ii) The set B is called the co-domain of the function f.

ii) The set of all elements in B which are the image of the elements of the domain A is called the range set of the function f range of $f = R(f) = \{y \in B \mid y = f(x) \text{ for some } x \in A\}$



1.3.3 Counting Number of functions from A to B :

Let $|A| = m$ & $|B| = n$, then the number of functions from A to B are $|B|^{|A|} = n^m$

Example :

Q. If $|A| = 3$ & $|B| = 5$, then find number of functions from A to B.

Solution :

If $|A| = m, |B| = n$ &

$F = \{f \mid f : A \to B\}$ then

$|F| = n^m$

Here $m = 3$ and $n = 5$

$\therefore |F| = n^m = 5^3 = 125$

Q. Let $|A| = m$ & $|B| = n$ defline
$F_1 = \{f | f : A \to B$ is a function$\}$ &
$F_2 = \{f | f : B \to A$ is a function$\}$. Then total number of functions form F_2 to F_1 are

(a) m^{n^m} (b) $n^{m^{n+1}}$
(c) $m^{n^{m+1}}$ (d) n^{m^n}


Solution: Here $|A| = m$, $|B| = n$
$\therefore |F_1| = |B|^{|A|} = n^m$
$|F_2| = |A|^{|B|} = m^n$
$F = \{f | f : F_2 \to F_1 \; be \; a \; function\}$
Then
$|F| = |F_1|^{|F_2|} = (n^m)^{(m^n)}$
$= n^{m \times m^n} = n^{m^{n+1}}$
$|F| = n^{m^{n+1}}$
$\therefore$ Number of functions from F_2 to F_1 are $n^{m^{n+1}}$

1.4 Classification of function:-

On the basis of mapping of elements, there are five types of functions
(1) one to one function (Injective function)
(2) Many to one function
(3) Onto function (Surjective function)
(4) Into function
(5) One-one & onto function (Bijective function or Invertible function)

1.4.1 One-one function (Injective function):

A function $f : A \to B$ is said to be one-one function, if two distinct elements of the domain have distinct images in the co-domain.
i.e. f is one-one iff for any $x_1, x_2 \in A$ if $x_1 \neq x_2 \Rightarrow f(x_1) \neq f(x_2)$
OR
If $f(x_1) = f(x_2) \Rightarrow x_1 = x_2$



Counting number of one-one function from A to B:
If $|A| = m$ & $|B| = n$. then total number of one-one functions from A to B are
$= \begin{cases} {}^n p_m & m \leq n \\ 0 & m > n \end{cases}$ where ${}^n p_m = \frac{n!}{(n-m)!}$

Example :-
Q. Let $A = \{1, 2, 3\}$ & $B = \{1, 2, 3, 4, 5\}$
suppose $F_1 = \{f | f : A \to$
B be function such that $f(1) = 1$ & $f(2) \neq 2$ $\}$ & $F_2 = \{f | f : A \to B$ be a one − one function such that $f(1) = 1$ & $f(2) \neq 2$ $\}$ then the cardinality of sets F_1 & F_2 respectively are ?

(1) 9, 9 (2) 20, 10
(3) 20, 9 (4) 18, 9

Solution:- Given $|A| = 3$ & $|B| = 5$
(i) Given $f(1) = 1$ & $f(2) \neq 2$
i.e. image of element 1 is fixed
$f(2)$ have 4 choices (i.e. 1, 3, 4, 5)
$f(3)$ have 5 choices (i.e. 1, 2, 3, 4, 5)
Hence, total number of functions in F_1 are $1 \times 4 \times 5 = 20$

(ii) Here $n = |B| = 5 > 3 = m = |A|$
$f(1) = 1$ & $f(2) \neq 2$
$\therefore$ $f(2)$ has three choices (i.e. 3,4,5)
& $f(3)$ has three choices of other then 1 and $f(2)$.
So,
Total number of functions in set F_2 is $1 \times 3 \times 3 = 9$
Hence, Option (3) is correct option

Q. The number of one-one function from set A to set B where $A = \{0, 1, 2, 3\}$ & $B = \{1, 2, 3, 4, 5\}$ are

(a) 60 (b) 40
(c) 100 (d) 120

Solution:- Here $|A| = m = 4$ and $|B| = n = 5$
$\Rightarrow m \leq n$
$\therefore$ Total number of one-one functions from A to B is ${}^n p_m$
$= {}^5 p_4 = \frac{5!}{(5-4)!} = \frac{5!}{1!} = 5! = 120$

1.4.2 Many-one function:

A function $f : A \to B$ is said to be many to one function, if there is at least two elements in the domain A has same image in B.

Note: A function that is not one-one is called to be many-one function.

Counting number of many to one function:
If $|A| = m$ & $|B| = n$ then
Total number of many-one function =
Total number of functions – Total number of one-one function
If $m > n$, then
Number of many to one function = $n^m - o = n^m$
If $m \leq n$, then
Number of many to one function = $n^m - {}^n P_m$

1.4.3 Onto (surjective) function:

A function $f: A \to B$ is said to be onto if each element of B has atleast one preimage in A.

Preimage: If f is function such that $y = f(x)$. Then we say
(i) y is image of x under f
(ii) x is preimage of y under f

Note: $f: A \to B$ is said to be onto if $\forall\, y \in B, \exists\, x \in A$ such that
$f(x) = y$

Counting number of onto functions form A to B:
1. If $|A| = m$ & $|B| = n$, then the number of onto functions from A to B is

$$= \begin{cases} n^m - \left[\sum_{r=1}^{n} (-1)^{r+1}\, n_{C_r} (n-r)^m\right], & m \geq n \\ 0 & m < n \end{cases}$$

Note:
(i) If range set is proper subset of B then f cannot be an onto function.
(ii) If $R(f) = B$, then f is onto

Examples:
The number of onto functions (surjective functions) from set $X = \{1, 2, 3, 4\}$ to set $Y = \{a, b, c\}$ is
(a) 36 (b) 64
(c) 81 (d) 72
Solution: $|X| = m = 4, |Y| = n = 3$
$\therefore$ total number of onto functions from X to Y
$= n^m - {}^nC_1(n-1)^m + {}^nC_2(n-2)^m - {}^nC_3(n-3)^m + \cdots$
$= 3^4 - {}^3C_1(3-1)^4 + {}^3C_2(3-2)^4 - {}^3C_3(3-3)^4 + \cdots$
$= 81 - 3(16) + 3(1)^4 = 81 - 48 + 3 = 36$
Total number of onto functions are 36
Hence, option (a) is correct.

1.4.4 Into function:

A function $f: A \to B$ is said to be into, if there exist an element in B which does not have preimage in A.
i.e. $f(A) \subsetneq B$

NOTE:
(i) A function which is not onto is an into function.
(ii) Number of into functions + number of onto functions = Total number of functions

Important Note:
(i) If $|A| = m$ & $|B| = n$, then total number of constant functions are n.
(ii) If $|A| = m$ & $|B| = 2$, then any function from A to B is into iff it is constant function.
$\Rightarrow$ Total number of into function from A to B is 2
Total number of functions from A to B is 2^m
$\therefore$ Total number of onto functions from A to B is $2^m - 2$.

1.4.5 One to one and onto OR Bijective OR invertible function:

A function $f: A \to B$ is bijective if it is one-one function and onto function.

Note:
(1) If $f: A \to B$ be a bijective function
$\Rightarrow$ (i) f is one-one $\Rightarrow |A| \leq |B|$
& (ii) f is onto $\Rightarrow |A| \geq |B|$
$\therefore$ If $f: A \to B$ be a bijective function then $|A| = |B|$

Converse need not be true.
i.e. if $|A| = |B| \not\Rightarrow f: A \to B$ is a bijective function

Example: If $f: [0, 1] \to [0, 1]$ such that $(x) = 1\ \forall x \in A$.
$\nexists$ any $x \in [0, 1]$ such that $f(x) = 0$
$\Rightarrow f$ is not onto function.
$\Rightarrow f$ is not a bijective function.

2. If $f: A \to B$ be a function where $|A| = |B| = n$ then f is one-one iff f is onto iff f is bijective
The above statement is not necessarily true when the cardinality of A & B are infinite.

Ex. $f: \mathbb{N} \to \mathbb{N}$ such that $f(n) = n + 1$
Here $|\mathbb{N}| = |\mathbb{N}|$ but f is not onto
$\because \nexists$ any $n \in \mathbb{N}$ such that $f(n) = 1$

Counting Number of bijective functions:
(i) If $|A| = |B| = n$, then number of bijective functions from A to B is equals to $n!$
(ii) If $|A| = |B| = n$, then number of functions from A to B which is neither one-one nor onto is equals to $n^n - n!$

Example:
If $A = \{1,2,3,4,5\}$ & $B = \{a, b, c, d, e\}$. Then
(a) The number of bijective functions
Answer: 5!
(b) The number of function which is one-on e but not onto
Answer: 0
(c) The number of function which is onto but not noe-one
Answer: 0
(d) The number of function which is neither one-one nor onto
$5^5 - 5!$


Note:
If $f: \mathbb{R} \to \mathbb{R}$ is a function then
(1) f is not one-one if graph of f cuts some line parallel to x –axis at least two points.
(2) f is one-one if graph of f cuts every lines parallel to x –axis at most one point.
(3) f is not onto if graph of f cuts some lines parallel to x –axis at no point.
(4) f is onto if graph of f cuts every lines parallel to x –axis at atleast one point.

Examples:
Q. If the functions $f, g: \mathbb{Z} \to \mathbb{Z}$ defined as $(n) = 3n + 2$ & $g(n) = n^2 - 5$, then
(1) Neither f nor g is one-one function
(2) The function f is one-one but not g

(3) The function g is one-one but not f
(4) Both f & g are one-one functions
Solution:-
(i) $f(n) = 3n + 2$
Let $n_1, n_2 \in \mathbb{Z}$ such that $f(n_1) = f(n_2)$
$\Rightarrow 3n_1 + 2 = 3n_2 + 2$
$\Rightarrow 3n_1 = 3n_2$
$\Rightarrow n_1 = n_2$
$\Rightarrow f$ is one-one function
(ii) $g(n) = n^2 - 5$
Here for any $n \in \mathbb{Z}$
$g(n) = n^2 - 5$ & $g(-n) = (-n)^2 - 5 = n^2 - 5$
But $n \neq -n$
$\Rightarrow g$ is not one-one function
Hence, option (2) is correct

Q. Let $f: X \to X$ such that $f(f(x)) = x$ for all $X \in X$. Then
(1) f is one-one and onto
(2) f is one-one but not onto
(3) f is onto but not one-one
(4) f need not be either one-one or onto
Solution:- for any $x \in X$, $f(f(x)) = x$
Let $x_1, x_2 \in X$ such that $f(x_1) = f(x_2)$
Apply f on above equation
$\Rightarrow f(f(x_1)) = f(f(x_2))$ $(\because f$ is a function$)$
$\Rightarrow x_1 = x_2$
$\Rightarrow f$ is one-one function
Onto :
(i) for any $x \in X$ (codomain), there is an element $f(x) \in X$ such that $f(f(x)) = x$
i.e. for any $i \in X$ (Codomain), $f(i)$ is a Preimage of i
Hence, f is onto function
OR,
If $f(g(x)) = x \quad \forall x \in X$ then f is onto & g is one-one function.
Consider $f(f(x)) = x$
$\Rightarrow f$ is onto & f is one-one function
Hence f is bijective map.
Thus, option (1) is correct.

Q. The function $f: \mathbb{R} \to \mathbb{R}$ defined by $f(x) = (x^2 + 1)^{2021}$ is
(1) Onto but not one-one
(2) One-one but not onto
(3) Both one-one & onto
(4) Neither one-one nor onto
Solution:- Here $f(x) = (x^2 + 1)^{2021}$
$f(-x) = ((-x)^2 + 1)^{2021}$ but $x \neq -x$
So, f is not one-one function
Consider $\forall x \in \mathbb{R}, x^2 + 1 \geq 1$
$\Rightarrow (x^2 + 1)^{2021} \geq (1)^{2021} = 1$
Hence



For any $y \in (-\infty, 1), \nexists$ any $x \in \mathbb{R}$ such that $f(x) = y$
Thus, f is not onto function.

Q. Let $\mathbb{Z}$ denote the set of integers and $\mathbb{Z}_{\geq 0}$ denotes the set $\{0, 1, 2, 3, \ldots.\}$. Consider the map $f: \mathbb{Z}_{\geq 0} \times \mathbb{Z} \to \mathbb{Z}$ Given by $f(m, n) = 2^m.(2n + 1)$. Then the map f is
(a) Onto (surjective) but not one-one (injective)
(b) One-one (injective) but not onto (surjective)
(c) Both one- one and onto
(d) Neither one-one nor onto
Solution :
Given, $f: \mathbb{Z}_{\geq 0} \times \mathbb{Z} \longrightarrow \mathbb{Z}$ defined by $f((m, n)) = 2^m(2n + 1)$
Let $(m_1, n_1), (m_2, n_2) \in \mathbb{Z}_{\geq 0} \times \mathbb{Z}$
Then, $f((m_1, n_1)) = f((m_2, n_2))$
$\Rightarrow 2^{m_1}(2n_1 + 1) = 2^{m_2}(2n_2 + 1)$
$\Rightarrow$ If $m_1 = m_2 \Rightarrow 2n_1 + 1 = 2n_2 + 1 \Rightarrow n_1 = n_2$
If $m_1 \neq m_2$, Then, without loss of generality we can consider that $m_1 > m_2$
$\Rightarrow 2^{m_1 - m_2}(2n_1 + 1) = (2n_2 + 1)$,
Which is a contradiction because term on left hand side is a multiple of 2 i.e even number and term on right hand side is an odd number.
So, $m_1 = m_2$ and $n_1 = n_2$
$\therefore f$ is an one-one function.
As, $0 \in \mathbb{Z}$ and $f(m, n) = 2^m(2n + 1) \neq 0 \; \forall (m, n) \in \mathbb{Z}_{\geq 0} \times \mathbb{Z}$
$(\because 2^m(2n + 1) = 0 \Rightarrow n = -\frac{1}{2} \notin \mathbb{Z})$
$\Rightarrow f$ is not an onto function.
Hence option (b) is correct

Q. Let $A = \{x^2 : 0 < x < 1\}$ and $B = \{x^3 : 1 < x < 2\}$ Which of the following statements is true?
(a) There is a one to one, onto function from A to B
(b) There is no one to one, onto function from A to B taking rational to rational.
(c) there is no one to one function from A to B which is onto
(d) There is no onto function from A to B Which is one to one

Solution: $A = \{x^2 : 0 < x < 1\}$ and $B = \{x^3 : 1 < x < 2\}$
A and B are open intervals and in metric space (R, d) open intervals are homeomorphic
So, there exists a homeomorphism of A onto B.
$\therefore$ there is one-one and onto function from A to B.
$\therefore$ option (a) is correct.

Q. How many onto function from set A = { 1,2,3,4,5,6,7,8,9,10} to the set B = { a, b }
(a) 1024 (b) 1022
(c) 100 (d) 98
Answer : (2)


Q. Let $f(n) = n^3 - 3n$ defined on set of integers is injective function. [T/F]
Answer : False

Note:
(1) If $f: A \to B$ is an invertible function then $f^{-1}: B \to A$ is also invertible function where $f^{-1}(y) = x$, whenever $f(x) = y$
(2) Let $f: A \to B$ is bijective function then
(i) $f \circ f^{-1} = I$ where $f \circ f^{-1}: B \to B$
(ii) $f^{-1} \circ f = I$ where $f^{-1} \circ f: A \to A$

Result:
1. If $f: X \to Y$ is one-one function & $A, B \subseteq X$, then
(i) $f(A \cup B) = f(A) \cup f(B)$
(ii) $f(A \cap B) = f(A) \cap f(B)$

2. If $f: X \to Y$ be a function such that f^{-1} exist & $A, B \subseteq Y$ then
(i) $f^{-1}(A \cup B) = f^{-1}(A) \cup f^{-1}(B)$
(ii) $f^{-1}(A \cap B) = f^{-1}(A) \cap f^{-1}(B)$
(iii) $f^{-1}(A^c) = \left(f^{-1}(A)\right)^c$

Example:-
1. If $f: X \to Y$ is a function & $A, B, C \subseteq X$. If $f(\phi) = \phi$ then which of the following is/are true?
(1) $f(A \cup B) = f(A) \cup f(B)$
(2) $f(A \cap B) = f(A) \cap f(B)$
(3) $f(A \cap B) \subseteq f(A) \cap f(B)$
(4) $f(A \backslash B) = f(A) \backslash (B)$
Solution:
(a) $A, B \subseteq A \cup B \Rightarrow f(A), f(B) \subseteq f(A \cup B) \Rightarrow f(A) \cup f(B) \subseteq f(A \cup B)$
$x \in A \cup B \Rightarrow f(x) \in f(A \cup B)$
Hence $f(A \cup B) \subseteq f(A) \cup f(B)$
$\Rightarrow f(A \cup B) = f(A) \cup f(B)$



(b)
Let $X = \{a, b, c, d\}$, $Y = \{1,4\}$ and let $f: X \to Y$ such that $f(a) = 1, f(b) = 1, f(c) = 4, f(d) = 4$
$A = \{a, c\} \Rightarrow f(A) = \{1,4\}$
$B = \{b, d\} \Rightarrow f(B) = \{1, 4\}$
$\Rightarrow f(A) \cap f(B) = \{1,4\}$
But $A \cap B = \phi$
$\Rightarrow f(A \cap B) = \phi$
$\Rightarrow f(A \cap B) \neq f(A) \cap f(B)$
So, Option (b) is incorrect.
If $x \in A \cap B \Rightarrow x \in A \,\&\, x \in B \Rightarrow f(x) \in f(A) \,\&\, f(x) \in f(B)$
$f(x) \in f(A \cap B)$
$\Rightarrow f(A \cap B) \subseteq f(A) \cap f(B)$
Hence, option (c) is correct

Let $X = \{a, b, c, d\}$, $Y = \{1,4\}$
$A = \{a, b\}$ and $B = \{c, d\}$ and let $f: X \to Y$ such that $f(x) = 1 \; \forall x \in \{a, b, c, d\}$
Then
$f(A) = \{1\}$ and $f(B) = \{1\}$
Therefore, $f(A \backslash B) = \boldsymbol{f}(\{a, b\}\{c, d\}) = \boldsymbol{f}(\{a, b\}) = \{\mathbf{1}\}$
And $f(A) \backslash (B) = \{1\} \backslash \{1\} = \phi$
Hence, option (d) is incorrect

1.4.6 Cantor's Theorem:
There is no onto map from any non-empty set to its power set.
Note:- There exist a one-one map from set A to $P(A)$
Since $f: A \to P(A)$ is not an onto map.
i.e. potential of power set of any set is more than set itself.
i.e. (potential of $P(\mathbb{N})$ is more than $\mathbb{N}$)
Remark:-
For any infinite set A, $|A| \ll |P(A)| \quad (i.e\ |\mathbb{Z}| < |P(\mathbb{Z})|, |\mathbb{Q}| < |P(\mathbb{Q})|, etc)$
$\therefore$ We need to define symbols to denote cardinality of infinite sets of different potential.
These symbols are called transfinite cardinals.
$|\mathbb{N}| = \aleph_0$ (aleph null, aleph nought)
$|P(\mathbb{N})| = c$ (continum)

1.5 Similar /Equivalent Sets:

1.5.1 Definition:
Two sets A & B are said to be equivalent or similar if there is one to one correspondence between their elements.
i.e. there exist a bijective map from A to B and it is denoted by $A \sim B$.